

Resilient Fields, Profitable Yields: The Promise of Intercropping Systems in a Changing Climate for Economic Gains

Shaik Rishitha¹, Santosh, D.T.^{2*}, Sumit Ray¹, Harun I. Gitari³, Masina Sairam¹, Sagar Lalichetti¹, Dinkar J. Gaikwad², Satya Prasad Nanda¹ and Sagar Maitra¹

¹M.S. Swaminathan School of Agriculture, Centurion University of Technology and Management, Paralakhemundi, Odisha, India

²Centre of Excellence for Smart Agriculture, Centurion University of Technology and Management, Paralakhemundi, Odisha, India

³Department of Agricultural Science and Technology, Kenyatta University, Nairobi, Kenya

*Corresponding author: santosh.dt@cutm.ac.in

Received: 12-05-2025

Revised: 28-08-2025

Accepted: 07-09-2025

ABSTRACT

Climate change fatally impacts agricultural production. To address this, adopting climate-resilient cropping systems is crucial, and intercropping presents a strong option. Intercropping, which involves growing two or more crops on the same land, offers a sustainable solution to the challenges of climate change and profitability. Its benefits are widely recognized. This study examines intercropping performance, economic viability, and climate resilience. Intercropping also improves soil health, erosion control, and carbon sequestration while increasing resource-use efficiency through complementary water, nutrient, and light utilization. It enhances management of environmental stresses, which is vital amid climate change. Economically, intercropping cuts input costs, especially when legumes contribute biological nitrogen, increases land-use efficiency as shown by higher land equivalent ratios, stabilizes production, and diversifies farm income. It adapts to various agroecological conditions. To promote adoption, the study underscores the need for supportive policy, innovation, and better extension services. Overall, intercropping is an economically viable, sustainable, climate-smart cropping system.

Keywords: Climate-resilience, Resource use efficiency, Soil health, Land Equivalent Ratio (LER), Sustainable agriculture, Economic feasibility

In recent times, agriculture is facing a tremendous crisis with issues like degradation of soil and water, shrinkage of fresh water and landmass for crop cultivation, and weather anomalies due to climate change (Mohanty *et al.* 2016; Zaman *et al.* 2017; Maitra *et al.* 2024a; Santosh *et al.* 2024a). In this context, to feed the growing world population, the direction of agriculture should be sustainable and climate-resilient (Sahu *et al.* 2024; Mukesh *et al.* 2024). There

are several agronomic approaches, and of them, the intercropping system can be considered as a suitable option to combat many problems because of

How to cite this article: Rishitha, S., Santosh, D.T., Ray, S., Gitari, H.I., Sairam, M., Lalichetti, S., Gaikwad, D.J., Nanda, S.P. and Maitra, S. (2025). Resilient Fields, Profitable Yields: The Promise of Intercropping Systems in a Changing Climate for Economic Gains. *Agro Economist - An International Journal*, 12(02): 55-67.

Source of Support: None; **Conflict of Interest:** None 

its spatio-temporal diversity and climate-resilience nature (Sairam *et al.* 2025a, b; Maitra and Gitari, 2020; Maitra *et al.* 2023). Intercropping can be described as the practice of growing two or more crops in each area at particular spatial (e.g., row, strip, mixed) and temporal contexts (Maitra *et al.* 2019, 2025a; Krishna *et al.* 2024; Ray *et al.* 2025a). This is one of the foundations of crop diversification strategies to enhance the effectiveness of using resources, including light, water, and nutrients, and retain or increase productivity (Chappa *et al.* 2022; Ray *et al.* 2025a, b). Intercropping systems are designed to capitalize on ecological interactions such as niche differentiation, facilitation, and competition minimization, and by combining complementary plant species to achieve optimal performance of the system (Sarkar *et al.* 2000; Maitra *et al.* 2001; Mirriam *et al.* 2023).

Recent studies have shown that intercropping has beneficial effects on biodiversity, plant-microbe interactions, and nutrient cycling, which increases the productivity of the system overall, especially in monoculture systems (Ray *et al.* 2024; Maitra *et al.* 2025b). For instance, legume-based intercropping systems in combination with cereals allow biological fixation of nitrogen, increasing the soil fertility and decreasing reliance on chemical fertilizers (Jensen *et al.* 2020; Maitra *et al.* 2023b). Moreover, intercropping leads to ecological intensification through the enhancement of the soil structure, the increase of the soil organic carbon and the enhancement of the biological control of pests (Altieri *et al.* 2015; Maitra, 2018; Santosh *et al.* 2022). The meta-analyses revealed that diverse cropping systems are highly beneficial to ecosystem services including pest control and pollination and also decrease the use of agrochemicals (Yousefi *et al.* 2024; Maheswari *et al.* 2025; Nori *et al.* 2025). Therefore, intercropping is not a new form of farming, but is also scientifically sound and sustainable cropping system with a lot of empirical evidence to support it.

Climate change is becoming more and more dominant on agricultural systems across the globe in the form of increased temperatures, unpredictable rainfall, extended droughts, and frequent abnormal weather conditions. Such alterations are highly problematic to traditional monoculture systems that are based on mono-crop cultures and consistent

climatic patterns (Gaikwad *et al.* 2022). Without proper adaptation measures, it is estimated that the global crop productivity may reduce by a maximum of 14 percent by 2050 (Aynalem *et al.* 2026). Monocropping systems are low-genetic and functional diversity systems and are therefore susceptible to environmental stresses due to a lack of genetic and functional diversity to buffer stresses (Hossain *et al.* 2021a, b; Ray *et al.* 2024). These systems become more vulnerable to pest attacks, disease outbreaks, and abiotic stress, and, consequently, the variability of yield and financial instability increases (Manasa *et al.* 2021). Rather, diverse cropping systems such as intercropping systems are emerging as climate-smart cropping systems (Maitra *et al.* 2026). In addition to improving soil water conservation and efficiency, intercropping systems create improved microclimates. These microclimates reduce heat stress and limit evaporation loss of soil moisture (Salakinkop *et al.* 2024). The use of legumes in cropping systems also increases nitrogen availability. This practice provides stability in yields during moisture fluctuations. Intercropping enables diversification, which spreads the risk of total crop failure among multiple crops. As a result, there is a more stable output in the face of climatic uncertainty (Raseduzzaman and Jensen, 2017). This risk buffering ability is especially important to smallholder farmers in climate-sensitive areas.

Intercropping is a synergistic method that combines the ecological sustainability and economic viability. It provides ecological benefits such as the increase in resilience of the system, the efficiency of resource-use, the increase of biodiversity, and the stabilization of agroecosystem processes (Moreira *et al.* 2024). Crops have complementary interactions, which increase nutrient uptake, and improves the wellbeing of the soil (Pramanick *et al.* 2023). Intercropping systems play a role in climate change as well by carbon sequestration and decreased greenhouse gas emissions because of reduced use of synthetic inputs (Maitra *et al.* 2024b). Intercropping becomes an important element of sustainable and regenerative agricultural systems because of these ecological benefits. Economically, intercropping has been found to increase the efficiency of land-use, commonly in terms of Land Equivalent Ratio (LER), which is commonly greater than unity, thus more productive than sole cropping designs. It also offers diversified




Fig. 1: Intercropping system as a climate-resilient and economically feasible technology for sustaining crop productivity

sources of income, lower input expenses, and improves profitability (Maitra *et al.* 2025a). In addition, intercropping increases economic resilience by diversifying risk, that is, by distributing risks among various crops, and, therefore, being less prone to market fluctuations and climatic shocks (Manasa *et al.* 2018; Maitra *et al.* 2024a). This is especially beneficial to small holder and resource-poor farmers. The idea is similar to climate-smart agriculture (CSA), which targets to boost productivity, resilience, and restricts greenhouse gas emissions (Fig. 1). It has been indicated that CSA practices such as intercropping have a significant positive effect on the farm income, productivity, and adaptive capacity in a changing climatic condition (Lalichetti *et al.* 2024).

Mechanisms for Climate Resilience

Through a mix of ecological processes that optimize resource acquisition, reduce interspecific competition, and maintain system stability under stress, intercropping improves climate resilience. The complementarities in resource utilization (water, nutrients, and light), facilitation among crops, and enhanced soil-plant-microbe relations are the main triggers of these mechanisms (Li *et al.* 2020). Consequently, intercropped systems can withstand climatic stress like drought, heat, and nutrient limitation much better than monocropping systems.

Resource Use Efficiency

Efficient resource use in intercropping systems relies on optimal use of water, nutrients, and light through complementing relationships between component crops, leading to greater productivity than monocropping (Dugassa, 2023). According to recent

meta-analyses, intercropping can significantly boost resource partitioning and facilitation mechanisms, as well as water-use efficiency (a 29% increase), making the system more adaptable to climatic stress (Liu *et al.* 2025).

Water-smart technology

Water-use efficiency (WUE) is one of the most significant processes in which intercropping can increase climate resilience. In intercropping systems, the crops are usually characterized by complementary root structure, i.e., deep-rooted, and shallow-rooted species, which enable them to tap into soil moisture at varying depths, and hence they do not compete intensely with each other regarding water (Santosh and Maitra, 2022). Such vertical stratification of roots helps in the optimal exploitation of the available water in the soil at the various soil horizons. As an example, the system to combine shallow-rooted crops, such as maize, and deep-rooted legumes, such as soybean or pigeon pea, can tap into deep and surface soil moisture concurrently, leading to better general intake of water (Santosh *et al.* 2024b). Compared to monocropping systems, these complementary rooting patterns have been shown to be able to boost water-use efficiency by 15–30%, particularly in semi-arid and drought-prone settings. Moreover, intercropping alters the microclimate of the soil by increasing canopy cover, decreasing the evaporation of soil, and increasing moisture retention of the soil (Santosh *et al.* 2025). Root plasticity and adaptation further increase water uptake during stressful conditions, and this makes crops productive even when there is a shortage of water. Accordingly, the adoption of crops that are associated with opposite root systems

can be described as a water-smart technology, which enables the intercropping system to mitigate drought stress and ensure the maintenance of yield stability in the circumstances of fluctuating climatic conditions (Santosh and Tiwari, 2017).

Efficient nutrient management

Intercropping can greatly enhance the efficiency of nutrient use, namely nitrogen (N), both biologically and ecologically. The nature of leguminous crops takes center stage in this process because it is capable of fixing nitrogen in the atmosphere (N_2) by a symbiotic relationship with rhizobia bacteria. In intercropping of cereals and legumes, legumes obtain most of their nitrogen through atmospheric fixation and thus do not rely heavily on soil nitrogen. This enables companion non-legume plants (e.g., cereals) to more efficiently utilize the available soil nitrogen and reduce competition and promote overall nitrogen-use efficiency (Hauggaard-Nielsen and Jensen, 2005; Maitra, 2020). In addition to this, some of the biologically fixed nitrogen is also passed on by the legumes to the related crops as root exudates, decomposition of plant materials, and microbial interactions, thus creating a richer soil nutrient pool. This enhances the fertility of the soil and helps to maintain crop productivity in the long run. Further, the production of chemical nitrogenous fertilizer releases a sizable quantity of greenhouse gas, and the adoption of a legume-based intercropping system, thus, can be considered as a ecofriendly cropping system.

Recent research also points out the fact that intercropping improves the levels of nitrogen cycling and microbial activity in the rhizosphere, which increases the availability of nutrients and uptake efficiency (Jena *et al.* 2020). The integration of legumes not only decreases the use of synthetic nitrogen fertilizers but also removes the environmental problems of nitrate leaching and greenhouse gases emissions. Also, intercropping systems exhibit the phenomenon of nutrient complementarity in which various crops utilize different nutrient pools or forms within the soil, maximizing total nutrient uptake (Maitra *et al.* 2001; Duchene *et al.* 2017). This addition is not just to nitrogen but also to other nutrients like phosphorus and micronutrients, and increases the productivity and sustainability of the system.

Soil Health Improvement and Conservation

One of the key components of climate-resilient agriculture is soil health, which encompasses the physical, chemical, and biological traits that preserve crop yield and ecosystem stability. Intercropping improves soil health by making soils more fertile, increasing the level of organic matter, and minimizing soil degradation, including erosion and loss of nutrients. The intercropping systems help in the long-term conservation and sustainability of the soil through continuous soil cover and diversification of the root systems in the changing climatic conditions.

Erosion Control

The greatest risk is the erosion of the soil and therefore affecting the sustainability of agriculture is when the climatic condition is characterized by heavy rainfall and high winds. Intercropping prevents soil erosion through constant vegetative coverage that keeps the soil surface free of the impact of rainfall and decreases the speed of the runoff (Maitra *et al.* 2021). The numerous crops reduce the soil particles' detachment and loss by ensuring that there is a dense canopy and ground cover. Moreover, a variety of root systems bind the soil particles with a better efficiency and increase the soil aggregation, which makes the soil more resistant to erosion (Bourke *et al.* 2021). Better infiltration and minimized surface runoff also play a part in conserving the soil. All these factors make intercropping a very effective practice of soil degradation prevention and an enhanced land productivity.

Carbon Sequestration

Carbon sequestration is another important ecosystem service that helps in the mitigation of climatic change, besides enhancing the fertility of soils. By boosting biomass generation and introducing various organic residues into the soil, intercropping regimes lead to better carbon sequestration (Yang *et al.* 2023). Increased root biomass and rhizodeposition under intercropping systems cause an enhancement in microbial activity, which causes stabilization of soil organic carbon (SOC) in various fractions (Amundson and Biardeau, 2018). Research has shown that intercropping systems have a much higher SOC level than monocultures because of constant organic

additions and better soil aggregation (Yang *et al.* 2023). This not only enhances the soil structure and nutrient levels but also converts agricultural soils into carbon reservoirs, which help in mitigating global climate change.

Enhanced Nutrient Availability

Legume-based intercropping system enhancing N and P availability through the biological nitrogen fixation and improved P solubility. Besides, it facilitates the cycling and bioavailability of essential macro- and micronutrients such as potassium (K), calcium (Ca), magnesium (Mg), zinc (Zn), iron (Fe), and boron (B) through root exudation, microbes dynamics, and enhanced soil organic matter (SOM). These nutrients play important roles for plant growth, productivity, and soil health, making legume intercropping an effective tool for addressing nutrient deficiencies. Intercropping legumes with cereals enhances K availability through root exudation of organic acids, releasing K from non-exchangeable reserves. Maize + legume intercropping system has shown a 15–20% increase in soil exchangeable K compared to monocropping (Wang *et al.* 2024). Deep-rooting legumes like pigeon pea access subsoil K and recycle them to the topsoil through leaf litter and root turnover. Maize and soybean intercropping enhances K uptake and nutrient accumulation in roots and green biomass by 20% with optimal K application (80:60 kg/ha) compared to no K application (Ahmed *et al.* 2020). Legume intercropping also improves Ca and Mg availability, essential for cell wall stability, enzymatic functions, and photosynthesis. Ca-rich leaf litter from legumes like cowpea and groundnut enriches soil Ca upon decomposition, benefiting associated non-legume crops like cereals. Legume root exudates solubilize Mg from soil minerals, improving its availability. Furthermore, legume intercropping enhances micronutrient bioavailability, such as Zn, Fe, and B, through rhizosphere interactions and microbial activity, promoting enzymatic functions and stress resistance.

Improved Soil Physical Properties

Legumes play a significant role in enhancing soil physical properties, including bulk density, saturated hydraulic conductivity, and the stability

of cracking clay. This structural improvement helps to reduce soil erosion. As anchor crops, legumes effectively enhance soil structure and water infiltration, particularly under semiarid and rainfed conditions, making them ideal for intercropping with fast-growing, shallow-rooted crops (Maitra *et al.* 2021). Water availability is a crucial factor affecting productivity in grain legume-based intercropping systems in the current context of the climate change. Enhancing water use efficiency (WUE) in these systems helps maximize resource use, particularly by conserving water through early foliage cover and increased leaf area (Maitra *et al.* 2024a). However, under limited water conditions, WUE in intercropping can be higher than in sole crop cultivation (Gou *et al.* 2017). Legume residues with narrow carbon to-nitrogen (C/N) ratios decompose quickly, boosting SOM and improving soil aggregate formation, while reducing bulk density. Legumes improve SOM, soil structure, cultivability, air circulation, water retention, and buffering capacity (Maitra *et al.* 2024a). The decomposition of SOM releases nutrients for plant growth and promotes better soil aggregation, stability, and resistance to erosion (Yadav *et al.* 2017).

Biotic Stress Management

Biotic stresses, such as pests and diseases, are significant limitations to crop yield, and climate change tends to increase their severity. Intercropping improves the natural resilience of agroecosystems by amplifying biodiversity and facilitating ecological interactions in the regulation of the population of pests and pathogens (Beillouin *et al.* 2021). These systems reduce the use of chemical control measures and will encourage sustainable management strategies to control pests and diseases.

Natural Pest Control

Intercropping enhances natural pest management through the diversification of plants and the formation of complex canopy coverages that hamper pest movement. A variety of crop species may conceal the presence of host plants, which makes them hard to find by pests. Moreover, intercropping offers habitat and alternative food supply to useful organisms, including predators and parasitoids, which contributes to biological

pest control. Abundance and diversity of natural enemies are useful in increasing the suppression of pests and decreasing crop damage (Hatt *et al.* 2018). Such environmentally friendly mode reduces the application of pesticides and is part of ecofriendly agriculture.

Disease Suppression

The host dilution effect is considered one of the key factors of disease management in intercropping systems, such that the presence of non-host plants decreases the density and connectivity of susceptible hosts, thus limiting the transmission of pathogens. The heterogeneity in space in intercropping systems breaks the disease transmission routes and lowers the growth of epidemics. Moreover, the intercropping systems result in increased microbial diversity in the soil that supports antagonistic relationships and inhibits soil-borne pathogens. The favourable microclimatic conditions, like increased air circulation and low humidity in mixed canopies, further suppress the growth of pathogens.

Economic Feasibility and Profitability

The economic viability is one of the major concerns that dictate the use of intercropping systems, especially for smallholder farmers with limited resources. Intercropping improves the profitability of farms by increasing resource-use efficiency, stabilization of yields in fluctuating climatic conditions, and less reliance on expensive external inputs. Besides the rise in overall productivity, diversified systems also offer different sources of income and mitigate the risks of the economy, which makes them more resilient and financially viable as compared to monocropping systems (Stomph *et al.* 2020).

Yield Stabilization

Stabilization of the yield during climatic unpredictability is one of the greatest economic benefits of intercropping. Diversification is a natural insurance policy whereby, if one crop failed because of drought, heat stress, or a pest attack, the other companion crops might still be successful, thus protecting overall farm production. The stability is due to variation in crop physiology, growth period,

and stress tolerance, which minimizes the risk of the occurrence of crop failure. Empirically, the intercropping systems have been found to be less variable in yield than monocultures, particularly in stress-prone environments (Li *et al.* 2021). This economic advantage is because such resilience guarantees a stable income and minimizes financial losses for farmers.

Land Equivalent Ratio (LER) is a popular measure that is used to assess the productivity and economic efficiency of intercropping systems. It can be described as the proportionate land area necessitated under pure cultivating to generate the identical yield as that produced in a system of intercropping (Mead and Wiley, 1981). When the LER exceeds 1.0, there is a yield benefit of intercropping as compared to monocropping. Many studies show that LER values in cereal-legume intercropping systems are between 1.1 and 1.5, which means that intercropping can yield 10-50 percent more per unit of area than sole cropping. The main benefit of this is linked with the complementary resource utilization, the decrease in competition, and the enhancement of ecological interactions between crops. The increased LER values are directly proportional to the increased land-use efficiency and economic returns, especially in areas where land is scarce. Therefore, LER is one of the important indicators of proving the economic excellence of intercropping systems. Similarly, there is a concept of the Area Time Equivalent Ratio (ATER), which considers the spatio-temporal efficiency of an intercropping system. The value of ATER higher than unity, is advantageous (Maitra *et al.* 2019). The efficient land use over time is always desirable in the mixed stand (Table 1).

Input Cost Reduction

Intercropping systems greatly lead to a decrease in the cost of input, especially the use of fertilizers and pesticides, hence increasing the net farm income. Legumes present in the intercropping systems make possible the biological fixing of nitrogen, which may provide a significant part of the nitrogen needed by the companion crops (Jensen *et al.* 2020). This saves on the synthetic nitrogen fertilizers and results to saving of costs and enhanced environmental sustainability. Moreover, in intercropping, the greater the biodiversity, the higher the level of

Table 1: Profitable intercropping system with LER and ATER values

Intercropping System	Row ratio	LER	ATER	Reference
Maize + cluster bean	2:3	1.9	1.40	Nandi <i>et al.</i> 2022
Pearl millet + groundnut	1:1	1.75	1.49	Priya <i>et al.</i> 2023
Maize + chickpea	2:2	2.02	1.37	Sain <i>et al.</i> 2023
	2:3	1.73	1.53	
Baby corn + green gram	3:3	1.55	1.47	Kumar <i>et al.</i> 2024
Maize + cowpea	2:3	1.41	1.28	Nori <i>et al.</i> 2025
	2:2	1.35	1.37	
Maize + groundnut	2:2	1.53	1.46	Ray <i>et al.</i> 2025a
Pigeon pea + field bean	1:2	1.43	1.19	Devika <i>et al.</i> 2025
Maize + green gram	1:4	1.66	1.50	Shafiqullah <i>et al.</i> 2026

Table 2: Economics of intercropping systems with higher profit and benefit-cost ratio

Intercropping System	Row ratio	Gross return (₹ ha ⁻¹)	Net profit (₹ ha ⁻¹)	B:C ratio	References
Sorghum + cowpea	2:1	98363	55597	2.30	Chaudhary <i>et al.</i> 2020
Maize + cluster bean	2:3	222584	155470	2.32	Nandi <i>et al.</i> 2022
Maize + groundnut	2:3	127742	83590	1.89	Panda <i>et al.</i> 2022
Groundnut + black gram	4:1	232099	152346	2.90	Harshavardhan <i>et al.</i> 2022
Maize + chickpea	1:2	127782	76877	1.51	Sain, <i>et al.</i> 2023
Pearl millet + groundnut	1:1	153620	99860	1.86	Priya <i>et al.</i> 2025
Maize + cowpea	2:2	146701	89161	1.55	Maheswari <i>et al.</i> 2025

natural control of pests, which decreases the use of chemical pesticides. Research has shown that the input costs can be decreased without the yield reduction in properly managed intercropping systems (Mwadalu *et al.* 2022; Wang *et al.* 2025). The reduced input needs increase profitability, as well as minimize risks in production that are related to changes in input prices, which makes intercropping a cost-effective practice in farming (Table 2).

Income Diversification

Intercropping gives the farmers a variety of products on the same land, such as food grains, pulses, fodder, and, in some cases, fuel or fibre crops. Such diversification increases the stability of incomes through the diversification of financial risks in various crops and market opportunities. An intercropping system is likely to have different crops with different periods of growth and harvesting, which enables farmers to earn money at different

times of the year. This is a steady and continuous flow of cash, which is especially useful in the case of smallholder farmers who depend on frequent income to meet their household expenditures. In addition, it is a more secure method against market fluctuations in prices because the farmers will not be reliant on a single commodity. Intercropping systems offer a way forward to better livelihoods of farmers by offering a sustainable trend in terms of ecological advantages and economic robustness in the face of shifting climatic and market conditions.

Barriers to Adoption and Implementation

Although intercropping systems have been documented to have positive environmental and economic impacts, their progress in most of the regions has been a low-adoption behavior as a result of a set of technical, economic, and institutional issues. The practical problems that farmers have to deal with are labor, knowledge, and financial capacity, which

do not allow them to switch to a diversified system despite relinquishing conventional monocropping (Keller *et al.* 2024; Jat *et al.* 2025). These obstacles are especially high among smallholder farmers, where resource constraints and the aversion to risk affect decision-making.

Management Complexity

Increased management complexity is one of the major obstacles to the adoption of intercropping. Intercropping, in contrast to monocropping, means that it is necessary to plan the mix of crops, their spatial layout, and the time of work, which makes the work more demanding and intensive in terms of management. The variability in crop maturity, height, and harvesting needs complicates harvesting and field operations. This poses a challenge to mechanization since the majority of farm equipment is geared towards uniform crop plots and not mixed cropping. Research indicates that inadequacy of appropriate equipment and failure to adapt to the mechanized systems greatly limit intercropping practices, especially in developing nations. Moreover, the lack of farm machinery and small land parcels also limits mechanization, which reduces the appeal of labour-intensive intercropping regimes in farmers with labour shortages (Srikanthnaik, 2024). Therefore, the added complexity of operations tends to deter mass adoption in favour of possible long-run rewards.

Technical Knowledge Gap

The lack of technical knowledge to design and administer effective systems is also a major obstacle to intercropping. Effective intercropping requires the choice of mutually compatible crop species, knowledge of their growth processes, and efficient planting time, spacing, and the management of nutrients. Nonetheless, most farmers do not have access to good extension services, training programs, and decision-support tools that can help them to adopt these systems. As it has been shown, the lack of technical knowledge and awareness are significant barrier to the implementation of diversified and climate-resistant agricultural techniques. Furthermore, intercropping systems are highly site-specific and require a site-specific understanding of crop interactions, soil, and climate. Farmers are

reluctant to try any new cropping methods because of the perceived risk and uncertainty caused by this complexity. The absence of standardized recommendations and low externalization of research results also increases the knowledge gap and reduces the rate of adoption (Maitra *et al.* 2023).

Policy Recommendations and Future Outlook

The intercropping systems should be widely adopted with the support of the policy framework, technological innovations, and institutional support. Although intercropping has been shown to have clear advantages in the context of resilience, productivity, and sustainability, the upscaling of the technique relies on enabling environments that minimize risks and increase farmer capacity. To make intercropping a regular practice of climate-smartness, agricultural policies in the future should combine economic incentives, digital technologies, and the platforms of knowledge exchange.

Technological Innovation

The technological innovation is a major motivation towards the optimization of the intercropping systems and the management complexities. Remote sensing, IoT-based monitoring, and decision-support systems are precision agriculture tools that allow farmers to manage spatial variability and of diversified cropping systems to optimize inputs. Simulation models of crop and data-driven methods can be used to determine the most appropriate combinations of crops, planting geometries, and sowing dates depending on the agroclimatic areas. The tools enhance the prediction of yield outcomes, efficiency in the use of resources, and climate resilience in various scenarios. The recent studies indicate that intercropping plus modern technologies may significantly increase yields and decrease the environmental footprints, which proves the possibility of integrating intercropping and modern technologies. Therefore, the combination of precision farming and intercropping can help reduce the disparity between the old knowledge and the scientific achievements.

Extension Services

The extension services are also important in encouraging the adoption of intercropping since

the systems are site-specific in the application of knowledge and management skills. The enhancement of local knowledge networks, such as farmer training, demonstration plots, or participatory research, can assist farmers to overcome the trial-and-error period of diversification. The nexus between researchers, policymakers, and farmers is necessary to make sure that scientific knowledge is converted into solutions that can be implemented in the field. Research has highlighted the importance of constant advisory services, capacity building, and knowledge-sharing platforms in terms of the successful implementation of intercropping. Additionally, the digital extension can be improved through digital tools like mobile-based advisory services and AI-driven decision systems, which will contribute to better information diffusion and real-time support of the farmers. Enhancement of extension systems will increase the likelihood of uncertainty reduction, increase adoption rates, and make intercropping systems sustainable in the long term.

CONCLUSION

A comprehensive and sustainable farming approach that successfully combines ecological resilience with financial viability is intercropping. Compared to traditional monocropping systems, intercropping systems maximize the use of natural resources like light, water, and nutrients by growing several crops in the same area and time. This results in increased productivity and stability. These systems' natural diversity improves soil health, encourages carbon sequestration, and lessens environmental degradation—all of which greatly aid in mitigating and adapting to climate change. Complementary root systems, biological nitrogen fixation, and effective resource partitioning are some of the mechanisms that underpin intercropping and help crops endure climate challenges, including heat, drought, and irregular rainfall. Continuous ground cover and a variety of biomass further promote soil conservation, lessen erosion, and enhance long-term soil fertility. By reducing reliance on chemical inputs and improving agroecosystem sustainability, intercropping also reinforces natural pest and disease control through ecological interactions.

Intercropping offers several benefits from an economic standpoint, including lower input costs, increased land-use efficiency, and yield stabilization.

The idea of diversification ensures more consistent farm revenue by acting as a natural buffer against market and climate fluctuations. The adaptability of intercropping across various agroecological circumstances is further demonstrated by systems including row, strip, relay intercropping, and agroforestry models, which present chances for multi-tier productivity and ongoing revenue creation. Nevertheless, a number of obstacles, including greater management complexity, fewer mechanization possibilities, knowledge gaps, and initial investment requirements, limit the use of intercropping systems despite their advantages. These obstacles demonstrate the necessity of focused interventions to help farmers make the shift from monocropping to diversified systems. In the future, coordinated initiatives including improved extension services, technology advancements, and policy support will be necessary for intercropping to scale successfully. Precision farming and crop modelling tools, as well as incentive-based policies that acknowledge environmental benefits, can improve system adoption and efficiency.

REFERENCES

- Ahmed, A., Aftab, S., Hussain, S., Nazir Cheema, H., Liu, W., Yang, F. and Yang, W. 2020. Nutrient accumulation and distribution assessment in response to potassium application under maize-soybean intercropping system. *Agronomy*, **10**(5): 725-742.
- Altieri, M.A., Nicholls, C.I., Henao, A. and Lana, M.A. 2015. Agroecology and the design of climate change-resilient farming systems. *Agronomy for Sustainable Development*, **35**(3): 869-890.
- Aynalem, M., Ayalew, Z. and Terefe, A.T. 2026. Impact of climate smart agricultural practices on technical efficiency among high value vegetable farming in Northwestern Ethiopia. *Cogent Food and Agriculture*, **12**(1): 2641276.
- Amundson, R. and Biardeau, L. 2018. Soil carbon sequestration is an elusive climate mitigation tool. *Proceedings of the National Academy of Sciences*, **115**(46): 11652-11656.
- Beillouin, D., Ben-Ari, T., Malézieux, E., Seufert, V. and Makowski, D. 2021. Positive but variable effects of crop diversification on biodiversity and ecosystem services. *Global Change Biology*, **27**(19): 4697-4710.
- Bourke, P.M., Evers, J.B., Bijma, P., van Apeldoorn, D.F., Smulders, M.J.M., Kuyper, T.W. and Bonnema, G. 2021. Breeding beyond monoculture: Putting the "intercrop" into crops. *Frontiers in Plant Science*, **12**: 734167.

- Chappa, L.R., Mugwe, J., Maitra, S. and Gitari, H.I. 2022. Current status, and prospects of improving sunflower production in Tanzania through intercropping with Sunn hemp. *International Journal of Bioresource Science*, **9**(1): 1-8.
- Chaudhary, R., Gupta, S.K., Singh, M. and Kohli, A. 2020. Effect of intercropping on growth, yield and profitability of sorghum, pearl millet and cowpea. *Journal of Pharmacognosy and Phytochemistry*, **9**(5S): 179-182.
- Devika, A.R., Rehman, H.M., Laxman, N. and Reddy, S. 2025. Intercropping Indices and Land Use Efficiency in Pigeon pea-based Cropping Systems in Eastern Dry Zone of Karnataka, India. *PlantArchives*, **25**(2): 2907-2912.
- Duchene, O., Vian, J.F. and Celette, F. 2017. Intercropping with legumes for agroecological cropping systems: Complementarity and facilitation processes and the importance of soil microorganisms: A review. *Agriculture, Ecosystems and Environment*, **240**: 148-161.
- Dugassa, M. 2023. Intercropping as a multiple advantage cropping system. *International Journal of Research in Agronomy*, **6**(1): 44-50.
- Gaikwad, D.J., Ubale, N.B., Pal, A., Singh, S., Ali, M.A. and Maitra, S. 2022. Abiotic stresses impact on major cereals and adaptation options-A review. *Research on Crops*, **23**(4): 896-915.
- Gou, F., Yin, W., Hong, Y., van der Werf, W., Chai, Q., Heerink, N., Martin K. and van Ittersum, M.K. 2017. On yield gaps and yield gains in intercropping: Opportunities for increasing grain production in northwest China. *Agricultural Systems*, **151**: 96-105.
- Harshavardhan, K., Sivakumar, S.S., Gunasekar John, J., Albert Alex, V. and Padmanathan P.K. 2025. Productivity of Groundnut Cum Blackgram based Intercropping System under Different Crop Ratios. *Indian Journal of Agricultural Research*, **59**(3): 389-395.
- Hatt, S., Boeraeve, F., Artru, S., Dufrêne, M. and Francis, F. 2018. Spatial diversification of agroecosystems to enhance biological control and other regulating services: An agroecological perspective. *Science of the Total Environment*, **621**: 600-611.
- Hauggaard-Nielsen, H. and Jensen, E. S. 2005. Facilitative root interactions in intercrops. *Plant and Soil*, **274**(1): 237-250.
- Hossain, A., Islam, M.T., Maitra, S., Majumder, D., Garai, S., Mondal, M., Ahmed, A., Roy, A., Skalicky, M., Brestic, M. and Islam, T. 2021a. Neglected and underutilized crop species: are they future smart crops in fighting poverty, hunger and malnutrition under changing climate. In Zargar et al. (eds) *Neglected and Underutilized Crops - Towards Nutritional Security and Sustainability*. Springer, Singapore. https://doi.org/10.1007/978-981-16-3876-3_1.
- Hossain, A., Pramanick, B., Bhutia, K.L., Ahmad, Z., Moulick, D., Maitra, S., Ahmad, A. and Aftab, T. 2021b. Emerging roles of osmoprotectant glycine betaine against salt-induced oxidative stress in plants: a major outlook of maize (*Zea mays* L.) p. 567-587. In T. Aftab et al. (eds.) *Frontiers in plant-soil interaction*. Academic Press. <https://doi.org/10.1016/B978-0-323-90943-3.00015-8>.
- Jat, R.K., Meena, V.S., Durgude, S., Sohane, R.K., Jha, R.K., Kumar, A. and Lenka, A.K. 2025. Bridging the gap: Challenges and adoption of climate-resilient agriculture technologies in agricultural landscapes across agro-climatic zones of Bihar, India. *Frontiers in Sustainable Food Systems*, **8**: 1504388.
- Jena, J., Palai, J.B. and Maitra, S., 2020. Modern concepts of fertilizer application. In: Maitra, S. and Pramanick, B. (eds) *Advanced Agriculture*, pp. 388-408.
- Jensen, E. S., Carlsson, G. and Hauggaard-Nielsen, H. 2020. Intercropping of grain legumes and cereals improves the use of soil N resources and reduces the requirement for synthetic fertilizer N: A global-scale analysis. *Agronomy for Sustainable Development*, **40**(1).
- Keller, C., Joshi, S., Joshi, T., Goldmann, E. and Riar, A. 2024. Challenges for crop diversification in cotton-based farming systems in India: A comprehensive gap analysis between practices and policies. *Frontiers in Agronomy*, **6**: 1370878.
- Krishna, T.G., Maitra, S., Kalasare, R.S., Ray, S., Mahto, R.K. and Sairam, M. 2024. Influence of location-specific nutrient management and plant stand on the growth, productivity, and nutrient uptake of irrigated *rabi* maize (*Zea mays* L.). *Research on Crops*, **25**(4): 560-569.
- Kumar, V., Ram, V. and Rajkumari, P. 2024. Evaluating intercropping indices, yield attributes and yield of baby corn and mungbean under different planting patterns. *International Journal of Plant and Soil Science*, **36**(8):460-469. <https://doi.org/10.9734/ijpss/2024/v36i84876>.
- Lalichetti S, Maitra S, Singh S, Masina SR. 2024. Advanced strategies for optimization of primary nutrients requirement in rice-A review. *Plant Science Today*, **11**(1): 353-365.
- Li, C., Hoffland, E., Kuyper, T.W., Yu, Y., Zhang, C., Li, H. and van der Werf, W. 2020. Syndromes of production in intercropping impact yield gains. *Nature Plants*, **6**(6): 653-660.
- Li, X.F., Wang, Z.G., Bao, X.G., Sun, J.H., Yang, S.C., Wang, P. and Li, L. 2021. Long-term increased grain yield and soil fertility from intercropping. *Nature Sustainability*, **4**(11): 943-950.
- Liu, H., Gao, X., Li, C., Cai, Y., Song, X. and Zhao, X. 2025. Intercropping increases plant water availability and water use efficiency: A synthesis. *Agriculture, Ecosystems and Environment*, **379**: 109360.
- Maheswari, N., Maitra, S., Sairam, M., Ray, S., Sagar, L., Santosh, D.T. and Gaikwad, D.J. 2025. Impact of real-time nitrogen management on the performance of maize-cowpea intercropping system. *Crop Research*, **60**(3,4): 139-146.

- Maitra, S. 2020. Intercropping of small millets for agricultural sustainability in drylands: A review. *Crop Research*, **55**(3,4): 162-71.
- Maitra, S. and Gitari, H.I. 2020. Scope for adoption of intercropping system in organic agriculture. *Indian Journal of Natural Science*, **11**(63): 28624 - 28631.
- Maitra, S. 2018. Role of intercropping system in agricultural sustainability. *Centurion Journal of multi discipline Research*, **8**(1): 77-90.
- Maitra, S., Ghosh, D.C., Sounda, G. and Jana, P.K. 2001. Performance of intercropping legumes in finger millet (*Eleusine coracana*) at varying fertility levels. *Indian Journal of Agronomy*, **46**(1):38-44.
- Maitra, S., Hossain, A., Brestic, M., Skalicky, M., Ondrisik, P., Gitari, H., Brahmachari, K., Shankar, T., Bhadra, P., Palai, J.B., Jena, J., Bhattacharya, U., Duvvada, S.K., Lalichetti, S. and Sairam, M. 2021. Intercropping – A Low Input Agricultural Strategy for Food and Environmental Security. *Agronomy*, **11**(2): 343.
- Maitra, S., Palai, J.B., Manasa, P. and Kumar, D. P. 2019. Potential of intercropping system in sustaining crop productivity. *International Journal of Agriculture, Environment and Biotechnology*, **12**(1): 39-45.
- Maitra, S., Patro, T.S.S.K., Reddy, A., Hossain, A., Pramanick, B., Brahmachari, K., Prasad, K., Santosh, D.T., Mandal, M., Shankar, T. and Sagar, L. 2023b. Brown top millet (*Brachiaria ramosa* L. Stapf; *Panicum ramosum* L.) – a neglected and smart crop in fighting against hunger and malnutrition, p. 221-245 In: M. Farooq *et al.* (eds.) *Neglected and underutilized crops*, Academic Press. <https://doi.org/10.1016/B978-0-323-90537-4.00012-0>.
- Maitra, S., Ray, S., Sagar, L., Sairam, M., Pramanick, B., Gitari, H. and Chappa, L. R. 2025a. Cropping system approach in climate-resilient food crop production. P. 327-356. In Pramanick, B. *et al.* (eds) *Climate-Smart Agricultural Technologies. Sustainability Sciences in Asia and Africa*. Springer, Singapore. https://doi.org/10.1007/978-981-96-7699-6_15.
- Maitra, S., Sahoo, U., Sairam, M., Gitari, H. I., Rezaei-Chiyaneh, E., Battaglia, M. L. and Hossain, A. 2023a. Cultivating sustainability: A comprehensive review on intercropping in a changing climate. *Research on Crops*, **24**(4): 702-715.
- Maitra, S., Sairam, M. and Santosh, D.T. 2024b. Evaluation of the growth, flower quality, yield and vase life of gerbera (*Gerbera jamesonii* L.) cultivars under open ventilated saw-tooth type polyhouse. *Research on Crops*, **25**(1): 104-109.
- Maitra, S., Sairam, M., Ray, S., Santosh, D.T., Gaikwad, D. J., Gitari, H. I. and Maheswari, N. 2024a. Applauding soil quality improvement through legume-based intercropping system: A review. *International Journal of Bioresource Science*, **11**(1): 59-69.
- Maitra, S., Sairam, M., Sagar, L., Ray, S., Santosh, D.T., Gaikwad, D.J., Atapattu, A.J. and Gitari, H.I. 2025b. Alley Cropping: A Sustainable Agroforestry System for Enhancing Productivity, Soil Health and Biodiversity. P:235-248. In: *Plant Biotechnology and Sustainable Agriculture: Bridging the Gap for Global Food Security*, <https://doi.org/10.1079/9781800627659.0016>.
- Maitra, S., Samui, S.K., Roy, D.K. and Mondal, A.K. 2001. Effect of cotton-based intercropping system under rainfed conditions in Sundarban Region of West Bengal. *Indian Agric*, **45**(3,4): 157-162.
- Maitra, S., Santosh, D.T., Jena, J., Behera, S.D., Ray, S., Sairam, M., Palai, J.B., Gaikwad, D. J., Rishitha, S., Gitari, H.I. and Atapattu, A.J. (2026). Understanding plant responses to abiotic stress through systems biology. p. 59-100 In: T, Aftab *et al.* (eds.) *Systems Biology for Crop Improvement: Emerging Tools and Technologies*. Academic Press. <https://doi.org/10.1016/B978-0-443-36457-0.00011-0>.
- Manasa, P., Maitra, S. and Reddy M.D. 2018. Effect of summer maize-legume intercropping system on growth, productivity and competitive ability of crops. *International Journal of Management, Technology and Engineering*, **8**(12): 2871-2875.
- Manasa, P., Sairam, M. and Maitra, S. 2021. Influence of maize-legume intercropping system on growth and productivity of crops. *International Journal of Bioresource Science*, **8**(1): 21-28.
- Mead, R. and Wiley, J. 1981. A review of statistical ideas relevant to intercropping research. *Journal of the Royal Statistical Society: Series A*, **144**(4): 462-487.
- Miriam, A., Mugwe, J., Raza, M.A., Seleiman, M.F., Maitra, S. and Gitari, H.H. 2022. Aggrandizing soybean yield, phosphorus use efficiency and economic returns under phosphatic fertilizer application and inoculation with *Bradyrhizobium*. *Journal of Soil Science and Plant Nutrition*, **22**(4): 5086-5098.
- Mohanty, S.S., Dash, N., Nanda, S.P. and Dey, R.K. 2016. Synthesis, characterization and removal of heavy metal ions from water bodies using novel hybrid materials bases on silica. *ISOR Journal of Applied Chemistry*, **9**(10): 2278-5736.
- Moreira, B., Gonçalves, A., Pinto, L., Prieto, M.A., Caroch, M., Caleja, C. and Barros, L. 2024. Intercropping systems: An opportunity for environment conservation within nut production. *Agriculture*, **14**(7): 1149.
- Mukesh, G., Sairam, M., Maitra, S., Gaikwad, D.J., Sagar, L. and Ray, S. 2024. Agrometeorological Indices, Physiological Growth Parameters and Performance of Finger Millet as Influenced by Different Cultivars Under Hot and Sub-Humid Region of Odisha. *International Journal of Experimental Research and Review*, **42**: 148-161.
- Mwadalu, R., Mochoge, B., Mwangi, M., Maitra, S. and Gitari, H.I. 2022. Response of Gadam sorghum (*Sorghum bicolor*) to farmyard manure and inorganic fertilizer application, *International Journal of Agriculture, Environment and Biotechnology*, **15**(1): 51 - 60.

- Nandi, S., Sairam, M., Panda, M., Shankar, T. and Maitra, S. 2022. Dry Matter Accumulation, Yield and Economics of Maize Cultivation as Influenced by Mixed Stands of Maize + Vegetable Legumes. *International Journal of Bioresource Science*, **9**(2): 93-100.
- Nori, M., Masina, S., Sagar, M., Sumit, R., Hemasree, K.R., Roman Kumar, M., Dinkar, J.G. and Rajesh Shriram, K., 2025. Influence of precision nitrogen management in Rabi maize and cow pea intercropping system under varied planting proportions in south Odisha. *Plant Science Today*, **12**(3).
- Panda, S. K., Sairam, M., Sahoo, U., Shankar, T. and Maitra, S. 2022. Growth, productivity and economics of maize as influenced by maize-legume intercropping system. *Farming and Management*, **7**(2): 61-66.
- Pramanick, B., Bera, A., Saha, P., Barman, A., Paramanick, B., Maitra, S. and Hossain, A. 2023. Soil microflora and their interaction with plants under changing climatic scenarios. P:16-40. In P. Mathur et al. (eds.) Microbial symbionts and plant health: trends and applications for changing climate Singapore: Springer Nature Singapore. https://doi.org/10.1007/978-981-99-0030-5_2.
- Priya, G.S., Maitra, S., Shankar, T. and Sairam, M. 2023. Effect of the summer pearl millet-groundnut intercropping system on the growth, productivity, and competitive ability of crops under south Odisha conditions. *Plant Science Today*, **10**(4): 238-246.
- Priya, G.S., Maitra, S., Masina, S. and Ray, S. 2025. Spatial arrangement in Pearl Millet-Groundnut intercropping system influences productivity, competition and economics. *International Journal of Experimental Research and Review*, **47**: 108-119.
- Raseduzzaman, M.D. and Jensen, E.S. 2017. Does intercropping enhance yield stability in arable crop production? A meta-analysis. *European Journal of Agronomy*, **91**: 25-33.
- Ray, S., Maitra, S., Sairam, M., Sameer, S., Sagar, L., Divya, B.S. and Gitari, H.I. 2025b. The nexus between intercropping systems, ecosystem services and sustainable agriculture: A review. *Research on Crops*, **26**(1): 1-11.
- Ray, S., Maitra, S., Sairam, M., Sravya, M., Priyadarshini, A., Shubhadarshi, S. and Padhi, D.P. 2024. An unravelled potential of foliar application of micro and beneficial nutrients in cereals for ensuring food and nutritional security. *International Journal of Experimental Research and Review*, **41**(Spl Vol): 19-42.
- Ray, S., Maitra, S., Sairam, M. and Sagar, L. 2025a. Impact of maize-groundnut intercropping system on growth and yield as influenced by phosphorus, potassium and sulphur fertilisation. *Crop Research*, **60**(5,6): 308-316.
- Sahu, S., Nanda, S.P. and Padhy, R.K. 2024. Interfacial Micro-heterogeneous Surface Phenomenon in the Oxidative Transformation of 2-Phenylethanol by Ce (IV). *Chemistry Africa*, **7**(3): 1605-1617.
- Sain, S., Maitra, S., Shankar, T. and Sairam, M. 2023. Competitive ability of legume-based intercrop for rabi maize (*Zea mays* L.) in north eastern ghats of Odisha. *Crop Research*, **58**(3,4): 136-143.
- Sairam, M., Maitra, S., Ray, S., Maity, B. and Pradhan, P. 2025a. Influence of need-based nutrient management on yield and nutrient use efficiency of Rabi maize (*Zea mays* L.) under sandy loam soils of Odisha, India. *Farming & Management*, **10**(1): 11-19.
- Salakinkop, S.R., Talekar, S.C., Patil, C.R., Patil, S.B., Jat, S.L., Iliger, K.S. and Kachapur, R.M. 2024. Sustainable intensification of climate-resilient maize-chickpea system in semi-arid tropics through assessing factor productivity. *Scientific Reports*, **14**(1): 3958.
- Santosh, D.T. and Maitra, S. 2022. Effect of drip irrigation and plastic mulch on yield and quality of ginger (*Zingiber officinale*). *Research on Crops*, **23**(1): 211-219.
- Santosh, D.T., Debnath, S., Maitra, S., Sairam, M., Sagar, L.L., Hossain, A. and Moulick, D. 2024a. Alleviation of climate catastrophe in agriculture through adoption of climate-smart technologies. P: 307-332 In: U, Chatterjee et al. (eds.) Climate Crisis: Adaptive Approaches and Sustainability. Springer Nature, Switzerland. https://doi.org/10.1007/978-3-031-44397-8_17.
- Santosh, D.T., Maitra, S., Sairam, M. and Gaikwad, D.J. 2024b. Onion growth, yield and water productivity as influenced by irrigation levels under drip irrigation and plastic mulch. *Research on Crops*, **25**(4): 626-634.
- Santosh, D.T., Pholane, L.P., Maitra, S., Mandal, D., Sairam, M., Gaikwad, D.J. and Ray, S. 2025. Effects of drip fertigation and plastic mulch on growth and productivity of okra. *Research on Crops*, **26**(3): 498-504.
- Santosh, D.T. and Tiwari, K.N. 2017. Response of tissue cultured banana (*Musa acuminate* L.) cv. Grand Naine to different levels of nutrients under drip fertigation and black plastic mulch. *Applied Ecology and Environmental Research*, **15**(4): 1473-1488.
- Santosh, D.T., Sain, S. and Maitra, S. 2022. Estimation of crop water requirement in maize and chickpea for different patterns of intercropping. *Indian Journal of Natural Sciences*, **13**(71): 41392-41398.
- Sarkar, R.K., Shit, D. and Maitra, S. 2000. Competition functions, productivity and economics of chickpea (*Cicer arietinum*)-based intercropping system under rainfed conditions of Bihar plateau. *Indian Journal of Agronomy*, **45**(4): 681-686.
- Shafiqullah, H., Asmatullah, D., Noorali, N. and Aminullah, N. 2026. Effect of Intercropping and Plant Geometry of Maize (*Zea mays* L.) Intercropped with Mungbean (*Vigna radiate* L.) on Growth and Yield Characters. *Archives of Current Research International*, **26**(3): 238-245.
- Srikanthnaik, J. 2024. Advancing agricultural mechanization in India: Challenges, opportunities, and farmer-centric solutions. *International Journal of Research in Agronomy*, **7**(6): 768-773.

- Stomph, T.J., Dordas, C., Baranger, A., de Rijk, J., Dong, B., Evers, J.B. and van der Werf, W. 2020. Designing intercrops for high yield, yield stability and efficient use of resources: Are there principles? P: 1-50. In: L,Donald(eds)Advances in Agronomy, Academic press, 160:1-50. <https://doi.org/10.1016/bs.agron.2019.10.002>.
- Wang, X., Li, Y., Yang, M., Liang, W. and Zhang, X. 2025. Intercropping with appropriate nitrogen reduction achieves the trade-off among soil biological health, soil multifunctionality, and crop productivity. *Field Crops Research*, **333**: 110085.
- Wang, J., Zhang, X., Wang, H., Liu, T., Fayyaz, A., Gonzalez, N.C., Wang, J., Chen, X., Zhao, J. and Yan, W.2024. Leguminous crop restores the carbon flow attenuation from nitrogen loading within soil nematode food web in a *Camellia oleifera* plantation. *Journal of Environmental Management*, **349**: 119580.
- Yadav, G.S., Lal, R., Meena, R.S., Babu, S., Das, A., Bhomik, S.N., Datta, M., Layak, J. and Saha, P. 2017. Conservation tillage and nutrient management effects on productivity and soil carbon sequestration under double cropping of rice in North Eastern Region of India. *Ecological Indicators*, **105**: 303-315.
- Yang, L., Luo, Y., Lu, B., Zhou, G., Chang, D., Gao, S. and Cao, W. 2023. Long-term maize and pea intercropping improved subsoil carbon storage while reduced greenhouse gas emissions. *Agriculture, Ecosystems and Environment*, **349**: 108444.
- Yousefi, M., Marja, R., Barmettler, E., Six, J., Dray, A. and Ghazoul, J. 2024. The effectiveness of intercropping and agri-environmental schemes on ecosystem service of biological pest control: A meta-analysis. *Agronomy for Sustainable Development*, **44**(2): 15.
- Zaman, A., Zaman, P. and Maitra, S. 2017. Water resource development and management for agricultural sustainability. *Journal of Applied and Advanced Research*, **2**(2): 73-77.

